


Geology - Interpreting the past to provide for the future



Prepared in cooperation with the
U.S. GEOLOGICAL SURVEY




MAP SERIES 66
Torrington 1:100,000 - scale
Preliminary Geologic Map



Base map from U.S. Geological Survey 1:100,000 - scale
metric topographic map of the Torrington, Wyoming -
Nebraska 30' x 60' minute Quadrangle, 1976
Projection: Universal Transverse Mercator (UTM), zone 13
North American Datum of 1983 (NAD83)
10,000-meter grid ticks: UTM, zone 12
10,000-foot grid ticks: Wyoming State Plane Coordinate
System, east zone and Nebraska north zone

UTM GRID CONVERGENCE (N)
1976 MATHEMATICAL DATUM AT CENTER OF SHEET
DISGROSS IS APPROXIMATE

WYOMING QUADRANGLE LOCATION



GEOLOGIC MAP OF THE TORRINGTON 30' x 60' QUADRANGLE, GOSHEN AND PLATTE COUNTIES, WYOMING, AND SIOUX AND SCOTTS BLUFF COUNTIES, NEBRASKA

compiled and mapped by

J. Fred McLaughlin and Ray E. Harris

2005

REFERENCES CITED AND SOURCES OF GEOLOGIC DATA

(numbers are those on INDEX TO GEOLOGIC MAPPING)

- Denson, N.M., and Botzly, T., 1949, Geology of the Hartville uplift, eastern Wyoming: U.S. Geological Survey Oil and Gas Investigations Preliminary Map 102, sheet 1, scale 1:50,000.
- Denson, N.M., 1974, Geologic map of the Lusk area, Goshen and Niobrara Counties, Wyoming: U.S. Geological Survey Open-File Map Series MS-83, plate 1, scale 1:125,000.
- Harris, R.E., McLaughlin, J.F., and Jones, R.W., 2008a, Geologic map of the Guernsey Quadrangle, Platte County, Wyoming: Wyoming State Geological Survey Map Series MS-83, scale 1:250,000.
- Harris, R.E., McLaughlin, J.F., and Jones, R.W., 2008b, Geologic map of the Guernsey Reservoir Quadrangle, Platte County, Wyoming: Wyoming State Geological Survey Map Series MS-83, scale 1:250,000.
- Lillegruen, J.A., 1993, Correlation of Paleozoic strata across Wyoming—a user's guide, in Snook, A.W., Steinhilber, J.E., and Roberts, S.B., editors, Geology of Wyoming: Wyoming State Geological Survey Memoir 5, v. 1, p. 414-477.
- Love, J.D., Christiansen, A.C., and Sever, C.K., 1980, Geologic map of the Torrington 1° by 2° Quadrangle, southeastern Wyoming and western Nebraska: U.S. Geological Survey Miscellaneous Field Studies Map MF-1184, scale 1:250,000.
- Love, J.D., Denson, N.M., and Botzly, T., 1949, Geology of the Glendo area: U.S. Geological Survey Oil and Gas Investigations Preliminary Map 102, sheet 1, scale 1:50,000.
- McGrew, L.W., 1963, Geologic map of the Laramie area, Platte and Goshen Counties, Wyoming: U.S. Geological Survey Bulletin 114141, 99, plate 1, scale 1:131,680.
- McLaughlin, J.F., 2008, Geologic map of the Douglas 30' x 60' Quadrangle, Converse and Platte Counties, Wyo.: Wyoming State Geological Survey Map Series MS-83, scale 1:100,000, 1 sheet, color.
- McGrew, L.W., 1967, Geologic map of the Antelope Gap Quadrangle, Platte County, Wyoming: U.S. Geological Survey Geologic Quadrangle Map GQ-603, scale 1:250,000.
- McGrew, L.W., 1967, Geologic map of the Chadron Hill Quadrangle, Goshen County, Wyoming: U.S. Geological Survey Geologic Quadrangle Map GQ-621, scale 1:250,000.
- McGrew, L.W., 1967, Geologic map of the Wheatland Quadrangle, Platte County, Wyoming: U.S. Geological Survey Geologic Quadrangle Map GQ-628, scale 1:250,000.
- McGrew, L.W., 1967, Geologic map of the Wheatland NE Quadrangle, Platte County, Wyoming: U.S. Geological Survey Geologic Quadrangle Map GQ-628, scale 1:250,000.
- Morris, D.A., and Babcock, H.M., 1960, Geology and ground-water resources of Platte County, Wyoming: U.S. Geological Survey Water-Supply Paper 1490, 195 p., plate 1, scale 1:125,000.
- Rapp, R.J., Visher, T.N., and Littleton, R.T., 1957, Geology and ground-water resources of Goshen County, Wyoming: U.S. Geological Survey Water-Supply Paper 1737, 145 p., plate 1, scale 1:125,000.
- Sando, W.J., and Sandberg, C.A., 1987, New interpretations of Paleozoic stratigraphy and history of the northern Laramie Range and vicinity, southern Wyoming: U.S. Geological Survey Professional Paper 1450, 79 p.
- Sims, P.K., Day, W.C., Snyder, G.L., and Wilcox, A.B., 1997, Geologic map of Precambrian rocks along part of the Hartville uplift, Platte and Goshen Counties, Wyoming: U.S. Geological Survey Miscellaneous Investigations Series Map 1:250,000, scale 1:250,000.
- McLaughlin, J.F., and Stafford, J.E., 2008, Preliminary geologic map of the Lusk 30' x 60' Quadrangle, Goshen and Niobrara Counties, Wyoming: Wyoming State Geological Survey Open File Report 08-5, scale 1:100,000, 1 sheet, color.
- Snyder, G.L., 1980, Map of Precambrian and adjacent Paleozoic rocks of the Hartville uplift, Goshen, Niobrara, and Platte Counties, Wyoming: U.S. Geological Survey Open-File Report 80-79, scale 1:100,000.
- Snyder, G.L., 1991, Hartville uplift, in Hanson, R.S., and others, editors, The Wyoming province, in Reed, J.C., Jr., and others, editors, Precambrian—contemporaneous U.S. Vol. C2 of The Geology of North America: Geological Society of America, Boulder, Colorado, p. 147-149.
- Snyder, G.L., and Peatman, Z.E., 1982, Precambrian geology and geochronology of the Hartville uplift, Wyoming: 1982 eastern geochronology field conference, p. 64-94.

Prepared in cooperation with and research supported by the U.S. Geological Survey, National Cooperative Geologic Mapping Program, under USGS award numbers 01HQAG0007, 01HQAG0045, and 01HQAG0144. The views and conclusions contained in this report are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

Digital cartography by J. Fred McLaughlin, Justin M. Mulhaly, Robin W. Lyons, Alice J. Vogelmann, Phyllis A. Ranz, and David W. Lucke

Map design and editing by Richard W. Jones

NOTICE TO USERS OF WYOMING STATE GEOLOGICAL SURVEY INFORMATION

Most information produced by the Wyoming State Geological Survey (WSGS) is public domain, is not copyrighted, and may be used without restriction. We ask that users credit the WSGS as a courtesy when using this information in whole or in part. This applies to published and unpublished materials in printed or electronic form. Contact the WSGS if you have any questions about citing materials or preparing acknowledgments. Your cooperation is appreciated.

The WSGS and the State of Wyoming disclaims any responsibility or liability for interpretations from this digital data or map printed from these digital data or decisions based thereon. The WSGS and the State of Wyoming retains and does not waive coverage immunity.

The use of or reference to trademarks, trade names, or other product or company names in this product is for descriptive or informational purposes, or is pursuant to licensing agreements between the WSGS or the State of Wyoming and software or hardware developers/vendors, and does not imply endorsement of those products by the WSGS or the State of Wyoming.

Users of these maps are cautioned about using the data at scales different than those at which the maps were compiled—using this data at a larger scale will not provide greater accuracy, and, in fact, it is a misuse of the data.

The Wyoming State Geological Survey (WSGS) and the State of Wyoming makes no representation or warranty, expressed or implied, regarding the use, accuracy, or completeness of the data presented herein, or from a map printed from these data, nor shall the act of distribution constitute any such warranty. The WSGS does not guarantee these digital data herein, or map printed from these data, to be free of errors or inaccuracies.

The WSGS and the State of Wyoming disclaims any responsibility or liability for interpretations from this digital data or map printed from these digital data or decisions based thereon. The WSGS and the State of Wyoming retains and does not waive coverage immunity.

The use of or reference to trademarks, trade names, or other product or company names in this product is for descriptive or informational purposes, or is pursuant to licensing agreements between the WSGS or the State of Wyoming and software or hardware developers/vendors, and does not imply endorsement of those products by the WSGS or the State of Wyoming.

Most information produced by the Wyoming State Geological Survey (WSGS) is public domain, is not copyrighted, and may be used without restriction. We ask that users credit the WSGS as a courtesy when using this information in whole or in part. This applies to published and unpublished materials in printed or electronic form. Contact the WSGS if you have any questions about citing materials or preparing acknowledgments. Your cooperation is appreciated.

(numbers are those listed in REFERENCES CITED AND SOURCES OF GEOLOGIC DATA)

EXPLANATION

DESCRIPTION OF MAP UNITS

Quaternary surficial deposits

Qal Alluvial deposits (Holocene)—Silt, fine-grained sand, and some gravel as valley fill in the present flood plains, bars, and islands of principal streams; includes gravel, sand, silt, mud, and clay deposited in and around reservoirs. Variable thickness; may exceed 200 feet (61 m) in Platte River valley (Rapp and others, 1957).

Qs Dune sand (Holocene)—Fine-grained, windblown sand; variable thickness

Qml Sand and loess (Holocene)—Fine-grained residual sand and loess; thickness from 0 to 40 feet (0 to 12 m)

Qsl Slope wash (Holocene and Pleistocene)—Pebbles, cobbles, and gravels amidst a variegated matrix; the result of mass wasting on steep slopes. Includes angular blocks of Paleozoic and Precambrian rocks at the base of the steeper slopes in the High Rock Range

Qml Mixed alluvium and colluvium (Holocene and Pleistocene)—Unconsolidated to poorly consolidated clay, silt, sand and gravel; mostly in flood plains and low terraces. Thickness generally 0 to 50 feet (0 to 15 m)

Ql Terrace gravels (Holocene and Pleistocene)—Pebble, cobble, and boulder deposits containing some silt, fine-grained sand, lenses of bentonitic clay, and locally some ash beds; thickness 0 to greater than 210 feet (64 m) on the highest terrace levels (Rapp and others, 1957)

Qsl Gravel deposits (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Qsl Gravels (Pleistocene)—Boulder to pebble conglomerate deposited in a fluvial environment by the ancestral North Platte River. Contains fluvial cross-bedded sand zones. The boulders include many types of crystalline, metamorphic, and sedimentary rocks including but not limited to granite, metarhyolite, amphibolite, schist, quartzite, marble, limestone, and sandstone. A fragment of black jade was found on the Guernsey Quadrangle (Harris and others, 2008a). Includes clasts from the Hartville uplift as well as clasts from the Laramie Mountains and farther west. Occurs as upland deposits in the Guernsey and Guernsey Reservoir areas (Harris and others, 2008a, 2008b) along the North Platte River. Thickness variable, but local thicknesses may exceed 150 feet (46 m) to as much as 225 feet (69 m) (Rapp and others, 1957)

Division 6 (Pennsylvanian and Upper Mississippian)—Well-indurated maroon to red orthoquartzite that firms cliffs and rocky knolls. Deposited on a well-developed karst surface, and fills sinkholes and covers in limestone of the underlying Guernsey Formation. Specially exposed and is absent in many areas within the quadrangle. This unit is equivalent to the Darwin Sandstone Member of the Hartville as described by Sando and Sandberg (1987). Thickness 0 to 120 feet (0 to 37 m)

Divisions 4, 5, and 6 undivided (Pennsylvanian and Upper Mississippian)

Guernsey Formation (Lower Mississippian and Upper Devonian)—Subdivided into an upper limestone unit of Early Mississippian age and a lower unit of Devonian age. The upper Guernsey consists of gray, cherty, coarsely crystalline, coarsely bedded, hard limestone; the chert has variety of colors and there are predominant dark-brown quartzite layers and nodules in this unit; average thickness approximately 135 feet (41 m). The surface of the upper unit is highly irregular, with many local variations (marked by a general thickening of ironstone toward the northwest). The lower unit is purple to gray dolomite, thin-bedded, silty, fine-grained, hard, brittle, and silty, interbedded with hard, dolomite purple shale and siltstone; thickness approximately 65 feet (20 m). Approximately 4 feet (1.2 m) of pink arkose with abundant feldspar and white quartz grains and pebbles occurs at base of lower unit and rests on Precambrian rocks. Total thickness exposed in the Hartville area is 140 to 260 feet (43 to 79 m). [Note: in places where this basal arkose and cross-bedded orthoquartzite occurs, it is mapped as Devonian or Cambrian quartzite (Q2) as described below.]

The basal arkose sandstone of the Guernsey was previously interpreted as Cambrian in age, but Sando and Sandberg (1987) reinterpreted this as the Fremont Canyon Sandstone (new name) and assigned it to the Upper Devonian. These authors considered the Guernsey Formation a superfluous name and divided it instead into the Madison Limestone (Mississippian in age) at the top, the Englewood Formation in the middle, and the Fremont Canyon Sandstone (Devonian in age) at the base.